

# Plasma phospholipid arachidonic acid content and calcium metabolism in idiopathic calcium nephrolithiasis

**BRUNO BAGGIO, ALESSANDRO BUDAKOVIC, MARIA ANGELA NASSUATO, GIUSEPPE VEZZOLI, ENZO MANZATO, GIOVANNI LUISETTO, and MARTINA ZANINOTTO**

*Division of Nephrology, Department of Medical and Surgical Sciences, University of Padua, Padua; Division of Nephrology Dialysis and Hypertension, San Raffaele Scientific Institute, Milan; and Central Laboratory, University-Hospital, Padua, Italy*

## **Plasma phospholipid arachidonic acid content and calcium metabolism in idiopathic calcium nephrolithiasis.**

**Background.** Reports of an increase in plasma and erythrocyte phospholipid arachidonic acid content and in urinary prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) excretion in patients with idiopathic calcium nephrolithiasis suggested their crucial role in the pathogenesis of hypercalciuria, a well-known risk factor for lithogenesis.

**Methods.** To confirm this hypothesis, 15 healthy subjects and 20 nephrolithiasis patients were evaluated for plasma phospholipid polyunsaturated fatty acid content and PGE<sub>2</sub> concentration, serum parathyroid hormone, 25 hydroxyvitamin D<sub>3</sub>, 1,25-dihydroxyvitamin D<sub>3</sub>, and bone-specific alkaline phosphatase levels, as well as urinary excretion of calcium, biochemical markers of bone resorption (hydroxyproline and crossLaps), and intestinal calcium absorption. Furthermore, the effect of a 30-day fish-oil diet supplementation on the previously mentioned parameters was investigated in the patients.

**Results.** At baseline, patients compared with controls showed higher levels of plasma phospholipid arachidonic acid content ( $P = 0.002$ ), PGE<sub>2</sub> ( $P = 0.0004$ ), serum 25-vitamin D<sub>3</sub> ( $P = 0.001$ ), and 1,25-vitamin D<sub>3</sub> ( $P = 0.001$ ), urinary excretion of calcium ( $P = 0.001$ ), hydroxyproline ( $P = 0.007$ ), and crossLaps ( $P = 0.019$ ), as well as intestinal calcium absorption ( $P = 0.03$  at 60 min). Fish oil supplementation induced a reduction in the plasma phospholipid arachidonic acid level ( $P < 0.0001$ ), and except for serum concentrations of 25-vitamin D<sub>3</sub>, normalized baseline blood and urinary parameters, including intestinal calcium absorption. A close correlation between plasma PGE<sub>2</sub> and serum 1,25-vitamin D<sub>3</sub> ( $P = 0.004$ ) and between phospholipid arachidonic acid and intestinal calcium absorption ( $P = 0.0002$ ) and calciuria ( $P = 0.007$ ) was observed, as well as between urine excretion of crossLaps and hydroxyproline ( $P < 0.0001$ ), crossLaps and calcium ( $P < 0.0001$ ), and hydroxyproline and calcium ( $P < 0.0001$ ).

**Conclusions.** These findings indicate that the phospholipid arachidonic acid content anomaly could represent the primary event responsible for the mosaic of metabolic and clinical alterations that are distinctive features of renal stone formers, and

suggest that a common pathogenetic mechanism might account for the several forms of hypercalciuria detected in idiopathic calcium nephrolithiasis.

We recently described an increased arachidonic acid (AA) level in the plasma and erythrocyte phospholipids of patients with idiopathic calcium nephrolithiasis (ICN) [1]. This finding followed previous literature reports of an alteration in urine excretion of prostaglandins E<sub>2</sub> (PGE<sub>2</sub>), the main metabolite of membrane phospholipid AA, in ICN patients [2–5]. In the light of the effects of dietary-induced phospholipid AA level modifications on cellular and renal calcium transport [6, 7] and the important hypercalciuric effect of PGE<sub>2</sub> [5, 8, 9], we advanced that phospholipid AA content might have a crucial role in the pathogenesis of hypercalciuria, a well-known risk factor for calcium lithogenesis [1, 10, 11]. To confirm this hypothesis, in a group of ICN patients, we evaluated the relationship between plasma phospholipid AA content, calcium-regulating hormones, and intestinal, renal, and bone calcium metabolism alterations, which were observed to constitute the main causes of idiopathic hypercalciuria in calcium nephrolithiasis [12].

## **METHODS**

### **Study patients**

The study was conducted in 20 recurrent idiopathic calcium oxalate stone formers (male 17 and female 3; median age 36 years; range, 23 to 56) consecutively selected among those attending our outpatient clinic and fulfilling the previously described criteria for ICN [1, 13]. Fifteen healthy staff members (male 12 and female 3; median age 34 years; range, 22 to 52) served as control subjects; none had a family or personal history of nephrolithiasis. All patients and control subjects had normal renal function, sterile urine samples, and normal blood pressure; none had diabetes mellitus or hepatic, thyroid,

**Key words:** calciotropic hormones, hypercalciuria, lithogenesis, renal stone formers, kidney stones.

Received for publication August 12, 1999

and in revised form January 26, 2000

Accepted for publication March 27, 2000

© 2000 by the International Society of Nephrology

or immunologic disease. All gave their informed consent to the study, according to the Helsinki Declaration. No patient or control subject had been placed on any type of therapeutic regimen in the eight weeks before the study, and in particular, none was taking any lipid-lowering drug or other agent known to affect lipid metabolism. In the eight weeks preceding the study, all participants were asked to consume a diet that matched their usual caloric content and macronutrient distribution. Seventeen of the patients agreed to continue the study and received one capsule (Seacor®-SPA-Italy) containing 850 mg of  $\omega$ -3 polyunsaturated fatty acid ethyl esters (fish oil) three times daily for 30 days, as previously reported [1].

### Analytical procedures

At the beginning of the study, all participants fasted from 8 p.m. of the preceding evening, and the following parameters were determined: creatinine, uric acid, urea nitrogen, electrolyte concentration, lipid analysis, and the phospholipid polyunsaturated fatty acid content [1], plasma PGE<sub>2</sub> (PerSeptive Diagnostics, Cambridge, MA, USA), serum bone-specific alkaline phosphatase (BAP; Boehringer Mannheim, Milan, Italy), intact parathyroid hormone (PTH; Bio-Rad Laboratories, Milan, Italy), 25-hydroxyvitamin D<sub>3</sub> (25-Vit D; Buhmann Laboratories, Allschwil, Switzerland), and 1,25-dihydroxyvitamin D<sub>3</sub> (1,25-Vit D; Nichols Institute, Diagnostics, San Juan Capistrano, CA, USA). Each analysis was carried out in duplicate and was expressed as the mean of two determinations. On the test day, after eight weeks of controlled diet, fasting urines were collected for two hours from 7 a.m. to 9 a.m. for assay of creatinine, urea nitrogen, calcium, and phosphorus, as well as some biochemical markers of bone turnover, including hydroxyproline (HP) (Beckman Analytical, Milan, Italy) and crossLaps (CTX; Cis Diagnostici, Vercelli, Italy). CTX measures the degradation products of type 1 collagen C-telopeptide and represents a specific marker of bone resorption [14]. All determinations were carried out in duplicate and were expressed as the mean for urine creatinine. Bone mineral density (BMD) of the lumbar spine (L2-L4) was assessed by dual-energy x-ray densitometer (Hologic QDR 1000, Boston, MA, USA). BMD values were expressed as the number of standard deviations from the mean of the normal young Caucasian population (T-score). Blood and urine parameters were re-evaluated at the end of the  $\omega$ -3 fatty acid supplementation trial.

### Intestinal strontium absorption test

The strontium (Sr) oral test was performed to determine intestinal calcium absorption as previously described [15] in 11 controls and in 17 patients before and after  $\omega$ -3 polyunsaturated fatty acid supplementation; no food was supplied during the Sr test. Sr absorption was expressed by two different methods. The first determined the incremental area under the serum concentration-time curve

(AUC at 30, 60, and 240 minutes) by the trapezoid method, and Sr absorption was expressed as mmol/L<sup>-1</sup>/minute; the other calculated the fractional absorption (FA) of the administered dose (FA at 30, 60, 240 minutes), considering 15% of body weight as the distribution volume [15, 16].

### Statistical analysis

Statistical evaluation was carried out with Student's *t*-test for paired and unpaired variables and determination of the *r* coefficient for linear correlation. The relationship between some parameters as dependent variables and others as independent variables was analyzed by a stepwise multiple regression model. Statistical significance was defined as a *P* value less than 0.05.

## RESULTS

Findings at baseline and after fish oil supplementation are summarized in Table 1. At baseline, the patients versus control subjects showed an increase in plasma phospholipid AA content and PGE<sub>2</sub> values and in serum 25-Vit D and 1,25-Vit D concentrations; all the other parameters, including age, sex ratio, body mass index (BMI), and plasma lipid values, did not differ between the two groups. At the urine level, the patients presented an increased excretion of calcium, CTX, and HP. Moreover, the patients showed an increase in intestinal absorption of Sr that was significant at 30 and 60 minutes (both expressed as FA and AUC) and at 240 minutes (expressed as AUC; Table 2). Mean T-scores in the two groups showed no significant difference; six patients had the T-scores that were a little lower, but not more than 10%.

The effects of fish oil supplementation are reported in Table 1. This regime induced a reduction in the plasma phospholipid content of linoleic and AA and an increase in eicosapentaenoic and docosahexaenoic acids. Moreover, the fish oil supplementation was able to modify the blood and urine parameters that were altered at baseline. It lowered calcium-regulating hormones, PGE<sub>2</sub> and 1,25-Vit D, but not 25-Vit D values, and reduced intestinal Sr absorption (Table 2), urine calcium excretion (Fig. 1), and the biochemical markers of bone remodeling, such as serum BAP and urine HP and CTX. No correlation was observed between the plasma phospholipid AA content and PGE<sub>2</sub> values (*r* = 0.14). In contrast, a direct significant correlation between plasma PGE<sub>2</sub> and serum 1,25-Vit D values emerged (Fig. 2), as well as a close correlation between the phospholipid AA level and intestinal Sr absorption (Table 3).

Moreover, a direct correlation between urine excretion of CTX and HP (Fig. 3), CTX and calcium (Fig. 4), and HP and calcium (Fig. 5), and between plasma phospholipid AA content and calciuria (Fig. 6) was observed. Stepwise multiple analysis showed that plasma phospho-

**Table 1.** Biochemical parameters of control subjects and patients before and after fish oil supplementation

	Controls	Patients		<i>P</i> values <sup>a</sup>	<i>P</i> values <sup>b</sup>	<i>P</i> values <sup>c</sup>
		Before	After			
Plasma fatty acid %						
C18:0 (Stearic)	13.50 ± 0.80	13.73 ± 1.29	14.18 ± 0.99	0.5474	0.0627	0.0406
C18:2 n-6 (Linoleic)	25.75 ± 2.55	24.67 ± 2.39	21.35 ± 3.02	0.2076	0.0001	0.0001
C20:4 n-6 (Arachidonic)	10.04 ± 0.91	11.17 ± 1.08	9.58 ± 1.17	0.0024	<0.0001	0.2336
C20:5 n-3 (EPA)	0.63 ± 0.16	0.77 ± 0.29	4.88 ± 0.88	0.1219	<0.0001	<0.0001
C22:6 n-3 (DHA)	3.82 ± 0.68	3.37 ± 0.94	6.44 ± 1.02	0.1177	<0.0001	<0.0001
Plasma phospholipids mg/dL	200.10 ± 25.10	201.40 ± 30.90	203.44 ± 28.02	0.8943	0.6656	0.7259
Plasma PGE <sub>2</sub> ng/L	111.73 ± 37.66	163.94 ± 39.24	95.82 ± 48.23	0.0004	<0.0001	0.3109
Serum PTH ng/L	32.93 ± 10.53	31.10 ± 11.35	37.41 ± 15.99	0.6290	0.0951	0.3636
Serum 25-vitamin D nmol/L	46.96 ± 11.71	62.55 ± 14.24	65.29 ± 20.27	0.0015	0.6038	0.0043
Serum 1,25 vitamin D pmol/L	44.45 ± 10.23	60.15 ± 15.58	44.33 ± 21.67	0.0018	0.0085	0.9848
Serum 25-vitamin D/1,25-vitamin D	1.15 ± 0.40	1.28 ± 0.53	2.06 ± 1.49	0.4355	0.0228	0.0291
Serum Calcium mmol/L	2.34 ± 0.07	2.29 ± 0.11	2.31 ± 0.09	0.1144	0.6049	0.3131
Serum Phosphorus mmol/L	1.29 ± 0.15	1.19 ± 0.18	1.22 ± 0.23	0.1155	0.6632	0.3300
Serum BAP U/L	28.10 ± 10.50	30.23 ± 16.34	26.94 ± 16.21	0.6623	0.0025	0.8146
Urinary urea mmol/mmol Cr	12.84 ± 2.45	13.91 ± 3.65	13.01 ± 4.00	0.3345	0.7088	0.8837
Urinary calcium mmol/mmol Cr	0.08 ± 0.03	0.13 ± 0.05	0.08 ± 0.04	0.0011	0.0348	0.9342
Urinary Phosphorus mmol/mmol Cr	0.35 ± 0.05	0.28 ± 0.06	0.31 ± 0.15	0.0017	0.1361	0.3802
Urinary CTX µg/mmol Cr	259.22 ± 57.76	323.82 ± 88.24	217.45 ± 96.58	0.0192	0.0179	0.1543
Urinary HP µmol/mmol Cr	12.00 ± 5.21	18.01 ± 6.96	11.12 ± 7.66	0.0078	0.0395	0.7102

Plus-minus values are means ± SD. Abbreviations are: P-Fatty acid, the percentage of fatty acids in plasma phospholipids; EPA and DHA, eicosapentaenoic and docosahexaenoic acids; BAP serum bone-specific alkaline phosphatase; CTX, crossLaps; HP, hydroxyproline.

<sup>a</sup> Comparison between patients (before) and control subjects (unpaired *t*-test)

<sup>b</sup> Comparison between patients before and after treatment (paired *t*-test)

<sup>c</sup> Comparison between patients (after) and control subjects (unpaired *t*-test)

**Table 2.** Strontium (Sr) oral load test in control subjects and patients before and after fish oil supplementation



	Controls	Patients		<i>P</i> values <sup>a</sup>	<i>P</i> values <sup>b</sup>	<i>P</i> values <sup>c</sup>
		Before	After			
Sr-FA 30 min	13.98 ± 6.14	18.92 ± 5.91	12.53 ± 5.29	0.0420	0.0002	0.5112
Sr-FA 60 min	22.84 ± 7.12	29.31 ± 8.00	24.75 ± 6.85	0.0380	0.0040	0.4836
Sr-FA 240 min	27.55 ± 10.72	30.87 ± 8.10	30.38 ± 10.38	0.3589	0.7951	0.4911
Sr-AUC 30 min	0.42 ± 0.19	0.57 ± 0.17	0.38 ± 0.16	0.0406	0.0002	0.5102
Sr-AUC 60 min	1.54 ± 0.53	2.03 ± 0.60	1.51 ± 0.48	0.0354	0.0001	0.8736
Sr-AUC 240 min	10.68 ± 3.57	12.95 ± 3.84	11.51 ± 3.29	0.1279	0.0182	0.5346

Plus-minus values are means ± SD. Sr-FA and Sr-AUC denote the intestinal Sr absorption expressed by two ways (**Methods**).

<sup>a</sup> Comparison between patients (before) and control subjects (unpaired *t*-test)


<sup>b</sup> Comparison between patients before and after treatment (paired *t*-test)

<sup>c</sup> Comparison between patients (after) and control subjects (unpaired *t*-test)

**Fig. 1.** Effects of fish oil administration on urine calcium excretion (*P* = 0.35).**Fig. 2.** Relationship between plasma prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) and serum 1,25-vitamin D<sub>3</sub> (1,25-Vit D) concentration in control subjects (○) and patients (●) (*r* = 0.47, *P* = 0.0044).

**Table 3.** Relationships between plasma phospholipid arachidonic acid content (P-AA) and intestinal strontium absorption rate expressed by fractional absorption and area under the serum concentration curve (AUC) procedures (**Methods** section)

	<i>r</i>	<i>P</i> values
P-AA/Sr-FA 30 min	0.41	0.0292
P-AA/Sr-FA 60 min	0.65	0.0002
P-AA/Sr-FA 240 min	0.66	0.0001
P-AA/Sr-AUC 30 min	0.41	0.0292
P-AA/Sr-AUC 60 min	0.54	0.0028
P-AA/Sr-AUC 240 min	0.68	0.0001



**Fig. 3.** Relationship between urine excretion of crossLaps (CTX) and hydroxyproline (HP) in control subjects (○) and patients (●) ( $r = 0.72$ ,  $P < 0.0001$ ).

lipid AA content and serum 1,25-Vit D concentration were the only statistically significant variables influencing intestinal Sr absorption at 60 minutes (Table 4), and that the calciuria was best explained by urine CTX and HP excretion, as well as intestinal calcium absorption (Table 5).

## DISCUSSION


The clinical and biochemical profiles of the patients who constituted the basis of our study showed the typical features of idiopathic calcium renal stone formers. Indeed, besides the hypercalciuria, we observed an alteration in calcium-regulating hormones, such as serum levels of calcidiol and calcitriol, and an increase in intestinal calcium absorption and bone turnover, as documented by the Sr test and the urinary excretion of biochemical markers of bone resorption (CTX and HP).

Hypercalciuria is the most common metabolic abnormality in ICN. It appears to reflect a heterogeneous entity and to result from a variety of causes, including dietary calcium excess, intestinal calcium hyperabsorption, calcium renal leak, and bone loss [12]. The problem is whether the several types of idiopathic hypercalciuria, including the absorptive, renal, and resorptive forms, reflect a spectrum of independent calcium disorders or

**Fig. 4.** Relationship between urine crossLaps (CTX) and calcium excretion in control subjects (○) and patients (●) ( $r = 0.75$ ,  $P < 0.0001$ ).**Fig. 5.** Relationship between urine HP and calcium excretion in subjects (○) and patients (●) ( $r = 0.72$ ,  $P < 0.0001$ ).

an epiphenomenon of a primary, as yet undefined generalized disturbance of calcium metabolism [17–19]. As 1,25-Vit D plays a central role in calcium metabolism at the intestinal, renal, and bone levels, a disorder in its pathway was a most attractive hypothesis to explain the different forms of hypercalciuria. Indeed, increased serum 1,25-Vit D levels have been observed in patients with hypercalciuria [17, 20–22] and have been confirmed by our study. Considering the selection criteria we used, the difference in 1,25-Vit D levels between patients and controls in our study did not depend upon different sunlight exposure times or other environmental factors. Moreover, calcitriol synthesis might be activated perhaps via a yet uncharacterized disorder in phosphate metabolism or, more simply, a mechanism(s) that up-regulates receptors for calcitriol. Both processes increase intestinal calcium absorption and urinary calcium excretion, and probably bone turnover as well [23–26]. However, most studies in this field have not yet reached definitive conclusions. In particular, it is not clear whether the disorder in the 1,25-Vit D pathway is a primary one or secondary to some other calcium metabolism defect. In our opinion,





**Fig. 6. Relationship between plasma phospholipid arachidonic acid (AA) content and calciuria in control subjects (○) and patients (●) ( $r = 0.44$ ,  $P = 0.0075$ ).**

**Table 4.** Multivariate regression analysis of the relationship between intestinal strontium absorption rate (Sr-FA 60 min) and other biochemical variables

Variables	Statistics $\beta$	SE	$P$ values
Sr-FA 60 min			
P-AA	0.666	(0.136)	0.0001
S-1,25 Vit D	0.351	(0.121)	0.0239
	$F = 14.440$	$R^2 = 0.53$	$P < 0.0001$

P-AA denotes the plasma phospholipid arachidonic acid content. The analysis was carried out in 28 subjects (11 controls and 17 patients) and is the result of a stepwise multiple regression by excluding the less significant variables.  $\beta$  denotes standardized regression coefficient.

the present study provides new and important insights into the pathophysiology of idiopathic hypercalciuria because it confirms our previous hypothesis of a causal link between membrane phospholipid AA levels and the calcium-related abnormalities frequently occurring in ICN patients [1, 10, 11]. This statement is supported by the results observed at baseline and is reinforced by the effects of polyunsaturated fatty acid administration. Indeed, we found an increase in the plasma phospholipid AA content, which agrees well with our previous finding in another group of nephrolithiasic patients [1]. As expected [1, 27], and also confirming good dietary compliance, fish oil supplementation was able to induce a reduction in the plasma phospholipid AA level. Moreover, it demonstrated a hypocalciuric effect that is in agreement with previous reports [1, 28]. This regimen also induced a reduction in  $PGE_2$  (the main metabolite of AA) and calcitriol values, as well as in intestinal Sr absorption, and a decrease in bone turnover, as documented by the reduction in biochemical markers of bone resorption (urinary HP and CTX excretion) and bone formation (serum BAP values).

Whatever the mechanism underlying the rise in the phospholipid AA proportion, we suggested that this anomaly could play a crucial role in the pathogenesis

**Table 5.** Multivariate regression analysis of the relationship between urine calcium excretion and other biochemical variables

Variables	Statistics $\beta$	SE	$P$ values
U-Ca			
U-CTX	0.452	(0.135)	0.0021
U-HP	0.407	(0.135)	0.0052
Sr-FA 60 min	0.265	(0.014)	0.0082
	$F = 27.844$	$R^2 = 0.73$	$P < 0.0001$

The analysis was carried out in 28 subjects (11 controls and 17 patients) and is the result of a stepwise multiple regression by excluding the less significant variables. Abbreviations are: CTX, crossLaps; HP, hydroxyproline; Sr-FA, 60 minutes of intestinal Sr absorption;  $\beta$ , standardized regression coefficient.

of idiopathic hypercalciuria through different actions. Indeed, an increase in phospholipid AA content determines a cascade of metabolic effects on calcium homeostasis leading to hypercalciuria, essentially because of a direct action of AA as second messenger on cellular calcium transport [29] or through the  $PGE_2/1,25$ -Vit D pathway.  $PGE_2$ , the main metabolite of AA, is known to modulate renal calcium handling by inhibiting the Na/K/2 Cl cotransporter, whose activity is important in driving calcium tubular reabsorption [5, 8] or by other mechanisms [9]. Moreover, the effects of  $PGE_2$  on the activity of 1,  $\alpha$  hydroxylase, the rate-limiting step in the formation of calcitriol, are well known [9], and seem to be confirmed by the direct correlation that emerged between  $PGE_2$  and 1,25-Vit D values (Fig. 1) and the results of fish oil supplementation, which showed a significant decrease in 1,25-Vit D levels, but not in those of its precursor 25-Vit D, leading to an increase in the calcidiol/calcitriol ratio. Excess calcitriol production was shown to induce intestinal calcium hyperabsorption, and reduced calcium tubular reabsorption [9]. Interestingly, a positive correlation was observed between plasma phospholipid AA level and intestinal Sr absorption (Table 3). Moreover, multiple regression analysis disclosed that the plasma phospholipid AA content and serum 1,25-Vit D levels were the only statistically significant and independent variables influencing intestinal Sr absorption (Table 4), thus suggesting a direct and independent effect of AA, besides the  $PGE_2/1,25$ -Vit D mechanism, on intestinal calcium transport.

Regarding the possible link between phospholipid AA values and bone remodeling, it is noteworthy that  $PGE_2$  and 1,25-Vit D, besides their effects on renal and intestinal calcium handling, also play a role in stimulating bone resorption and could constitute an additional source of urinary calcium in ICN [30]. The observations of primary or secondary high bone turnover in some hypercalciuric renal stone formers were confirmed by the present data. Indeed, we found an increase in specific biochemical markers of bone resorption, a positive correlation between urine excretion of calcium, CTX, and HP (Figs. 3 and 4), and a simultaneous reversal effect of dietary fish

oil administration on calciotropic hormones, biochemical markers of bone turnover, and urine calcium excretion. Calcitriol is a potent stimulator of osteoclast formation in vitro as well in vivo [31–33], while PGE<sub>2</sub> is an important factor of bone resorption directly or by the activation of 1,25-Vit D synthesis [34, 35]. On the other hand, it was reported that calcitriol may induce interleukin-1 (IL-1) osteoblastic receptor expression [36], and AA can modulate the synthesis of cytokines and growth factors [37, 38], well-known important local factors that are also involved in bone remodeling [39–41]. IL-1 and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) are among the most powerful stimulators of bone resorption and are well-recognized inhibitors of bone formation. The notion of a link between AA and bone turnover seems to be confirmed by previous findings that dietary fish oil supplementation reduced bone turnover in an experimental animal model [42] and decreased IL-1, IL-6, and TNF- $\alpha$  synthesis by circulating monocytes [40, 43]. Moreover, preliminary results from our laboratory demonstrated a specific effect of AA on the genetic expression of IL-1, IL-6, and TNF- $\alpha$  in a human osteoblastic cell line [44].

On the whole, the results of the present study seem to confirm the hypothesis of a close association in ICN patients between plasma phospholipid AA levels, calcium-regulating hormones, and urinary calcium excretion. A logical interpretation of our observations is that a higher phospholipid AA content would induce an increased calciotropic hormone concentration, thus leading to an increased intestinal calcium absorption, a decreased renal tubular calcium reabsorption, and an increased calcium bone loss, all mechanisms that have been observed to contribute to hypercalciuria in ICN. This idea is supported by multiple regression analysis that showed that urinary calcium excretion was well explained by the urine HP and CTX excretion and by intestinal calcium absorption (Table 5), thus suggesting that calciuria ultimately reflects the net rates of either intestinal calcium absorption or bone resorption, processes that are integrated by the calciotropic hormones, and perhaps other systems. In this setting, it is interesting to note that urinary calcium excretion was positively correlated with the plasma phospholipid AA level (Fig. 5).

One could speculate whether the phospholipid AA level anomaly really can explain the mosaic of metabolic and clinical alterations that are distinctive features of ICN patients, such as serum PGE<sub>2</sub> and 1,25-Vit D levels, kidney, intestine, and bone calcium metabolism, and several forms of idiopathic hypercalciuria [10]. In our opinion, further investigations in a larger number of hypercalciuric patients, classified according to the conventional criteria for differentiating hypercalciuria, will be important for recognizing in this anomaly, the primary event responsible for idiopathic hypercalciuria. Nonetheless, the results of the present study seem to reinforce our opinion and

that of others [17, 18, 45] that, despite differences in the clinical presentation, a common pathogenetic mechanism may account for the several subtypes of idiopathic hypercalciuria in ICN, which represents an epiphenomenon of a systemic disturbance in calcium metabolism.

Finally, in addition to the investigations of the yet unknown origin of the anomalous phospholipid AA level, to which genetic and nutritional/environmental factors seem to contribute [1], the positive and long-term effects of polyunsaturated fatty acid administration on the metabolic and clinical features of ICN patients and the natural history of this renal disease need further clinical verification. In this regard, it is worth noting that Eskimos, who eat a  $\omega$ -3 fatty acid-rich diet and seem to have a very low desaturase activity [46], the rate-limiting step in the biosynthetic pathway of highly unsaturated fatty acids [47], do not generally develop nephrolithiasis. Moreover, in studies of Japanese stone formers, it was found that the prevalence of nephrolithiasis since the Second World War correlated positively with the increase not only in protein, but also in animal fat [48, 49]. On the other hand, it is known that ICN emerged as a disease entity in Western countries at the beginning of this century, with an expanding societal weight running parallel with affluency.

## ACKNOWLEDGMENTS

We gratefully acknowledge Ms. Raffaella Marin for technical assistance and Ms. Patricia Segato for help in preparing the manuscript.

Reprint requests to Bruno Baggio, M.D., Dipartimento di Scienze Mediche e Chirurgiche, Policlinico Universitario, Via Giustiniani 2, 35120 Padova, Italy.  
E-mail: BBaggio@ux1.unipd.it

## REFERENCES

- BAGGIO B, GAMBARO G, ZAMBON S, MARCHINI F, BASSI A, BORDIN L, CLARI G, MANZATO E: Anomalous phospholipid n-6 polyunsaturated fatty acid composition in idiopathic calcium nephrolithiasis. *J Am Soc Nephrol* 7:613–620, 1996
- BUCK AC, LOTE CJ, SAMPSON WF: The influence of renal prostaglandins on urinary calcium excretion in idiopathic urolithiasis. *J Urol* 139:421–426, 1983
- HIRAYAMA H, IKEGAMI K, SHIMOMURA T, SOEJIMA H, YAMAMOTO T: The possible role of prostaglandins E<sub>2</sub> in urinary stone formation. *J Urol* 139:549–551, 1988
- HENRIQUES-LA ROCHE C, RODRIGUEZ-ITURBE B, HERRERA J, PARRA G: Increased urinary excretion of prostaglandin E<sub>2</sub> in patients with idiopathic hypercalciuria is a primary phenomenon. *Clin Sci* 75:581–587, 1988
- GAMBARO GF, MARCHINI MA, NASSUATO A, NARDELLOTTI S, CHECCHETTO L, CALÒ B, BAGGIO B: The hypercalciuric effect of PGE<sub>2</sub> in nephrolithiasis patients is mediated by inhibition of Na-K-2Cl cotransport activity, in *Renal Stones: Aspects of Their Formation, Removal and Prevention*, edited by TISELIUS HG, AKADEMITYCK AB, Sweden, Edsbruk, 1995, pp 95–96
- GAMBARO G, BORDONI A, HRELIA S, BORDIN L, BIAGI PL, SEMPLICINI A, CLARI G, MANZATO E, BAGGIO B: Dietary manipulation of delta-6-desaturase modifies phospholipid arachidonic acid levels, and the urinary excretion of calcium and oxalate in the rat: Insight in calcium lithogenesis *J Lab Clin Med* 135:89–95, 2000
- GAMBARO G, CHECCHETTO S, BACELLE L, BONVICINI P, BUDAKOVIC

- A, BAGGIO B: Dietary polyunsaturated fatty acids and idiopathic calcium nephrolithiasis, in *Proceedings of the 7th European Symposium on Urolithiasis*, edited by JUNGERS P, DAUDON M, Paris, Elsevier, 1997, p 88
8. BAGGIO B, GAMBARO G, MARCHINI F, VINCENTI M, CELOOTTO G, PESSINA A, SEMPLICINI A: Abnormal erythrocyte and renal frusemide-sensitive sodium transport in idiopathic calcium nephrolithiasis. *Clin Sci* 86:239–243, 1994
9. KUROKAWA K: Calcium-regulating hormones and the kidney. *Kidney Int* 32:760–771, 1987
10. BAGGIO B, PLEBANI M, GAMBARO G: Pathogenesis of idiopathic calcium nephrolithiasis: Update 1997. *Crit Rev Clin Lab Sci* 35:153–187, 1998
11. BAGGIO B, GAMBARO G: Abnormal arachidonic acid content of membrane phospholipids—the unifying hypothesis for the genesis of hypercalciuria and hyperoxaluria in idiopathic calcium nephrolithiasis. *Nephrol Dial Transplant* 14:553–555, 1999
12. PACK KC, KAPLAN R, BONE H, TOWNSEND J, WATERS O: A simple test for the diagnosis of absorptive, resorptive and renal hypercalciurias. *N Engl J Med* 292:497–500, 1975
13. BAGGIO B, GAMBARO G, MARCHINI F, CICERELLO E, TENCONI R, CLEMENTI M, BORSATTI A: An inheritable anomaly of red-cell oxalate transport in “primary” calcium nephrolithiasis correctable with diuretics. *N Engl J Med* 314:599–604, 1986
14. BONDE M, QUIST P, FLEDELIUS C, RIIS BJ, CHRISTIANSEN C: Immunoassay for quantifying type I collagen degradation products in urine evaluated. *Clin Chem* 40:2022–2025, 1994
15. VEZZOLI G, BARAGETTI I, ZERBI S, CAUMO A, SOLDATI L, BELLINZONI P, CENTEMERO A, RUBINACCI A, MORO GL, BIANCHI G: Strontium absorption and excretion in normocalciuric subjects: Relation to calcium metabolism. *Clin Chem* 44:586–590, 1998
16. MILSON S, IBBERTSON K, HANNAN S, SHAW D, PYBUS J: Simple test of intestinal calcium absorption measured by stable strontium. *BMJ* 295:231–234, 1987
17. COE FL, FAVUS MJ, CROCKETT T, STRAUSS AL, PARKS JH, PORAT A, GANTT CL, SHERWOOD LM: Effects of low-calcium diet on urine calcium excretion, parathyroid function and serum 1,25(OH)<sub>2</sub>D<sub>3</sub> levels in patients with idiopathic hypercalciuria and in normal subjects. *Am J Med* 72:25–32, 1982
18. COE FL, PARKS JH: Idiopathic hypercalciuria: The contribution of Dr. Jacob Lemann, Jr. *J Am Soc Nephrol* 5:S59–S69, 1994
19. LEVY F, ADAMS-HUET B, PAK C: Ambulatory evaluation of nephrolithiasis: An update of a 1980 protocol. *Am J Med* 98:50–59, 1995
20. KAPLAN RA, HAUSSLER MR, DEFTOS LJ, BONE H, PAK C: The role of 1 alpha, 25 dihydroxyvitamin D in the mediation of intestinal hyperabsorption of calcium in primary hyperparathyroidism and absorptive hypercalciuria. *J Clin Invest* 59:756–760, 1977
21. SHEN F, BAYLINK D, NIELSEN R, SHERRARD D, IVEY J, HAUSSLER M: Increased serum 1,25-dihydroxyvitamin D in idiopathic hypercalciuria. *J Lab Clin Med* 90:955–962, 1977
22. GRAY R, WILZ D, CALDAS A, LEMANN JJ: The importance of phosphate in regulating plasma 1,25-(OH)<sub>2</sub>-vitamin D levels in humans: Studies in healthy subjects in calcium-stone formers and in patients with primary hyperparathyroidism. *J Clin Endocrinol Metab* 45:299–306, 1977
23. INSOGNA K, BROADUS A, DREYER B, ELLISON A, GERTNER J: Elevated production rate of 1,25-dihydroxyvitamin D in patients with absorptive hypercalciuria. *J Clin Endocrinol Metab* 61:490–495, 1985
24. BROADUS AE, INSOGNA KL, LANG R, ELLISON AF, DREYER BE: Evidence for disordered control of 1,25-dihydroxyvitamin D production in absorptive hypercalciuria. *N Engl J Med* 311:73–80, 1984
25. BATAILLE P, ACHARD JM, FOURNIER A, BOUDAILLIEZ B, WESTEEL PF, ESPER EN, BERGOT C, JANS I, LALAU JD, PETIT J, HENON G, LAVAL JEANTET MA, BOUILLON R, SEBERT JL: Diet, vitamin D and vertebral mineral density in hypercalciuric calcium stone formers. *Kidney Int* 39:1193–1205, 1991
26. HESS B, CASEZ JP, TAKKINEN R, ACKERMANN D, JAEGER P: Relative hypoparathyroidism and calcitriol up-regulation in hypercalciuric calcium renal stone formers: Impact of nutrition. *Am J Nephrol* 13:18–26, 1993
27. FISHER S: Dietary polyunsaturated fatty acids and eicosanoid formation in humans. *Adv Lipid Res* 23:169–198, 1989
28. BUCK AC, DAVIES RL, HARRISON T: The protective role of eicosapentaenoic acid (EPA) in the pathogenesis of nephrolithiasis. *J Urol* 146:188–194, 1991
29. NAOR Z: Is arachidonic acid a second messenger in signal transduction? *Mol Cell Endocrinol* 80:C181–C186, 1991
30. WEISINGER JR: New insights into the pathogenesis of idiopathic hypercalciuria: The role of bone. *Kidney Int* 49:1507–1518, 1996
31. HOLTROP ME, COX KA, CLARK MR, HOLIOCK MF, ANST CS: 1,25-dihydroxy-cholecalciferol stimulates osteoclasts in rat bones in the absence of parathyroid hormone. *Endocrinology* 108:2293–2301, 1981
32. MCSHEEHY PM, CHAMBERS TJ: 1,25-dihydroxyvitamin D<sub>3</sub> stimulates rat osteoblastic cells to release a soluble factor that increases osteoblastic bone resorption. *J Clin Invest* 80:425–429, 1987
33. MAIERHOFER WJ, GRAY RW, CHEUNG HS, LEMANN JJ: Bone resorption stimulated by elevated serum 1,25(OH)<sub>2</sub> vitamin D concentrations in healthy man. *Kidney Int* 24:555–566, 1983
34. FILIPPONI P, MANNARELLI C, PACIFICI R: Evidence for a prostaglandin-mediated bone resorptive mechanism in subjects with fasting hypercalciuria. *Calcif Tissue Int* 43:61–66, 1988
35. NORRDMAN RW, JEE SS, HIGH WB: The role of prostaglandins in bone in vivo. *Prostaglandins Leukot* 41:139–149, 1990
36. LACEY DL, GROSSO LE, MOSER SA, ERDMANN J, TAN HL, PACIFICI R, VILLAREAL DT: IL-1 induced murine osteoblast IL-6 production is mediated by the type IL-1 receptor and is increased by 1,25 dihydroxyvitamin D<sub>3</sub>. *J Clin Invest* 91:1731–1742, 1993
37. SHERMAN ML, WEBER BL, DAITA R, KUFU DW: Transcriptional and posttranscriptional regulation of macrophage specific colony stimulation factor gene expression by tumor necrosis factor: Involvement of arachidonic acid metabolites. *J Clin Invest* 85:445–449, 1990
38. CUGHEY GE, MANTZIORIS E, GIBSON RA, CLELAND LG, JAMES MJ: The effect on human tumor factor alpha and interleukin 1 beta production of diets enriched in ω-3 fatty acids from vegetable oil or fish oil. *Am J Clin Nutr* 63:116–122, 1996
39. MANOLAGAS SC, JILKA RL: Bone marrow, cytokines, and bone remodeling. *N Engl J Med* 332:305–311, 1995
40. WEISENGER JR, ALONZO E, BELLORIN-FONT E, BLASINI AM, RODRIGUEZ MA, PAZ-MARTINEZ V, MARTINIS R: Possible role of cytokines on the bone mineral loss in idiopathic hypercalciuria. *Kidney Int* 49:244–250, 1996
41. PACIFICI R: Idiopathic hypercalciuria and osteoporosis—distinct clinical manifestations of increased cytokine-induced bone resorption? *J Clin Endocrinol Metab* 82:29–31, 1997
42. CLAASSEN N, POTGIETER HC, SEPPA M: Supplemented gamma-linolenic acid and eicosapentaenoic acid influence bone status in young male rats: Effects of free urinary collagen cross-links, total urinary hydroxyproline, and bone calcium content. *Bone* 16:385S–392S, 1995
43. PACIFICI R, RIFERS L, TEITLBAUM S, SLATOPOLSKY E, MILLER R: Spontaneous release of interleukin 1 from human blood monocytes reflects bone resorption in idiopathic osteoporosis. *Proc Natl Acad Sci USA* 84:4616–4620, 1987
44. PRIANTE G, BORDIN L, BRUNATI AM, ANGLANI F, CLARI G, BAGGIO B: Specific modulatory effect of arachidonic acid on bone cytochrome mRNA expression. *Ital J Miner Electrol Metab* (in press)
45. HEILBERG IP, MARTINI LA, DRAIBE SA, AJZEN H, RAMOS OL, SHOR N: Sensitivity to calcium intake in calcium stone forming patients. *Nephron* 73:145–153, 1996
46. ENGELMANN B, OP DEN KAMP JAF, ROELOFSEN B: Replacement of molecular species of phosphatidylcholine: Influence on erythrocyte Na transport. *Am J Physiol* 258:C682–C691, 1990
47. BRENNER RR: Factors influencing fatty acid chain elongation and desaturation, in *The Role of Fats in Human Nutrition* (2nd ed), edited by VERGROESEN AJ, CRAWFORD M, London, Academic Press, 1989, pp 45–79
48. IGUCHI M, KATAOKA K, KOHRI K, YACHIKU S, KURITA T: Nutritional risk factors in calcium stone disease in Japan. *Urol Int* 39:32–35, 1984
49. MASAI M, ITO H, KOTAKE T: The effect of dietary intake on urinary oxalate excretion in calcium renal stone formers. *Br J Urol* 76:692–696, 1995